



ASPIRIN USE

The following is a partial list of medications that can increase your bleeding tendency at the time of surgery and in the immediate post operative period.

This is a **PARTIAL** list. Any and all medications (including herbal products and vitamins) that you are taking must be reported to our office a minimum of **2 weeks prior to your surgery**. If you are unsure about a product you are using that is not on this list, call our office and ask the nursing staff. Check the labels of ALL medications that you take for ingredients that may be on this list.

DO NOT take any drugs on this list **2 weeks (3 weeks for any Diet Aids)** prior to or following your surgery without Dr. Caridi's consent. **If you are unsure of any product you wish to use, please call the office before taking it or use Tylenol.**

Aspirin	Bufferin	Diurex	Medipren	Sine Aid
Actifed	Buffex	Doan's Pills	Midol	Sudafed
Advil	Buf Tabs	Dolobid	Mobigesic	Supac
Aleve	Brufen	Dirstan	Monacet	Synalogs
Allegra D	Bupropion	Duradyne	Motrin	Talwin
Alka Seltzer	Cama	Easprin	Norgesic	Toradol
Aluprin	Cephalgesic	Ecotrin	Nardil	Triaminiein
Alcohol	Claritin D	Emprin	Nuprin	Trandate
Anacin	Clinoril	Emprazil	Nyquil	Trental
Anaprox	Congespirin	Excedrin	Oxycodone	Trilisate
Anaproxn	Cope	Feldene	Pabirin	Vanquish
Ansaid	Coricidan D	Fiorinal	Panalgesic	Voltaren
APA	Coricidin	Flurbiprophen	Pepto Bismol	Wesprin
Arthritis Pain	Tablets	Sodium	Percodan	Zactin
Reliever	Contact	Four Way	Persisrin	Zorpin
ASA	Coumadin	Cold Tabs	Quagesic	Zyrtec D
Asperbuf	Concerta	Ibuprofen	Rhinex	
Asriptin	Darvon with	Indocin	Robaxisal	
Aspergum	Aspirin	Indomethacin	Robaxin	
BC Powder	Diet Aids	Lodinc	Rufen	
Bayer	Decongestants	Meclomen	Sine Off	